Author Index

Aburatani, R., see Okamoto, Y. 367
Andersson, S.
——, Hedsten, U. and Jacobsson, S.
Quantitation of polydimethylsiloxane in pharmaceutical formulations by gel permeation chromatography 474
Anthony, L. J.
—— and Holland, R. A.
New method for surface characterization by gas chromatography 291
Balmér, K.
——, Persson, B.-A. and Schill, G.
Optimization of detection sensitivity for enantiomers of metoprolol on silica-bonded α_1-acid glycoprotein 107
Bartle, K. D., see Payne, K. M. 161
Bayer, E., see Koppenhoefer, B. 139
Bertsch, W., see Roberts, D. 59
Betina, V.
Chromatographic methods as tools in the field of mycotoxins (review) 187
Boshoven, C., see Tock, P. P. H. 95
Brinkman, U. A. Th., see Thiebaut, D. 151
Brownstein, A. D., see Sandra, P. 63
Burford, R. P., see Harahap, N. 53
Camilleri, P.
——, Dyke, C. and Hossner, F.
Chiral separation of the optical isomers of the antimalarial drug halofantrine 471
Chervet, J.-P., see Thiebaut, D. 151
Cook, C. E., see Lee, Y. W. 434
Cramers, C., see Hetem, M. 3
——, see Van Es, A. 39
Cruz, L. A.
—— and Jenke, D. R.
Evaluation of assay specificity in non-suppressed ion chromatography 271
Dalga, R. J.
—— and Lambert, W. J.
Development of a polymer-based reversed-phase high-performance liquid chromatographic stability indicating assay for U-78 608, an iron complexing monocarbam antibiotic 427
David, F., see Sandra, P. 63
Davies, I. L., see Payne, K. M. 161
Decottignies, P., see Le Maréchal, P. 305
De Haan, J., see Hetem, M. 3
De Jong, G. J., see Thiebaut, D. 151
Doehl, J.
—— and Greibrokk, T.
Identification of reaction products from the pyridinium dichromate derivatization of prostaglandins by high-performance liquid chromatography and direct chemical ionization mass spectrometry 345
Drevin, I.
——, Larsson, L. and Johansson, B.-L.
Column performance of Q-Sepharose HP in analytical- and preparative-scale chromatography 337
Dube, S., see Vohra, R. M. 463
Dyke, C., see Camilleri, P. 471
Elizarov, S. M.
Application of fast protein liquid chromatography for the isolation of vertebrate casein kinase-1 448
Engelhardt, H.
——, Gross, A., Mertens, R. and Petersen, M.
High-performance liquid chromatographic columns and stationary phases in supercritical-fluid chromatography 169
Érchegyi, J., see Vadász, Z. 377
Es, A. van, see Van Es, A. 39
Fang, Q. C., see Lee, Y. W. 434
Frei, R. W., see Thiebaut, D. 151
Gehas, J.
—— and Wetlaufer, D. B.
Hydrophobic interaction chromatography of small molecules: characterization of the retention of acyl coenzyme A homologues 249
Greibrokk, T., see Doehl, J. 345
Grob, K.
Concurrent eluent evaporation with co-solvent trapping for on-line reversed-phase liquid chromatography–gas chromatography. Optimization of conditions 73
Gross, A., see Engelhardt, H. 169
Gueldner, R. C., see Snook, M. E. 439
Guiochon, G., see Katti, A. M. 119
Guoshi, T., see Tun, L. 458
Guthrie, J. L., see Hull, C. J. 387
Haan, J. de, see Hetem, M. 3
Haken, J. K., see Harahap, N. 53
Harada, K.-I., see Ikai, Y. 397
Harahap, N.
——, Burford, R. P. and Haken, J. K.
Gas chromatographic analysis of thermoplastic aromatic polyamides after alkali fusion 53
Hatada, K., see Okamoto, Y. 367
Hedsten, U., see Andersson, S. 474

Henry, M.
——, Pauthe-Dayde, D. and Rochd, M.
Extraction and high-performance liquid chromatographic determination of gypsogenin 3,O-glucuronide 413
Herbst, H., see Hummel, M. 420
Hetem, M.
——, Rutten, G., Vermeer, B., Rijks, J., Van de Ven, L., De Haan, J. and Cramers, C.
Deactivation with polymethylhydrosiloxane. A comparative study with capillary gas chromatography and solid-state ^{29}Si nuclear magnetic resonance spectroscopy 3
Holland, R. A., see Anthony, L. J. 291
Hossner, F., see Camilleri, P. 471
Hull, C. J.
——, Guthrie, J. L., McKinney, R. W., Taylor, D. C., Mabud, Md. A. and Prescott, S. R.
Determination of toluenediamines in polyurethane foams by high-performance liquid chromatography with electrochemical detection 387
Hummel, M.
——, Herbst, H. and Stein, H.
Reversed-phase high-performance liquid chromatography of very long oligodeoxyribonucleotides 420
——, see Koppenhoefer, B. 139
Ikai, Y.
——, Oka, H., Kawamura, N., Yamada, M., Harada, K.-I., Suzuki, M. and Nakazawa, H.
Improvement of chemical analysis of antibiotics. XVI. Simple and rapid determination of residual pyridonecarboxylic acid antibacterials in fish using a prepacked amino cartridge 397
Ishii, D., see Takeuchi, T. 147
Ito, Y., see Lee, Y. W. 434
Jacobsson, S., see Andersson, S. 474
Jacquot, J.-P., see Le Maréchal, P. 305
Jenke, D. R., see Cruz, L. A. 271
Johansson, B.-L., see Drevin, I. 337
Jong, G. J. de, see Thiebaut, D. 151
Kaida, Y., see Okamoto, Y. 367
Kaiser, R. E.
—— and Rieder, R.
High boiling organic traces in drinking water. Quantitative analysis by liquid–liquid enrichment within the analytical glass capillary 49
Kakimoto, T., see Nishi, H. 259
Katti, A. M.
——, Ramsey, R. and Guiochon, G.
Quantitative extraction of thymine–thymine dimer from a large excess of thymine by preparative liquid chromatography 119
Kawai, S.
——, Nagano, H. and Maji, T.
Improved method for the separation of methylolmelamines by high-performance liquid chromatography 467
Kawamura, N., see Ikai, Y. 397
Kehl, M.
—— and Lottspeich, F.
Limitations of high-speed reversed-phase high-performance liquid chromatography observed with integral membrane proteins 131
Kołodziejczyk, A. M., see Wodecki, Z. J. 454
Koppenhoefer, B.
——, Muschalek, V., Hummel, M. and Bayer, E.
Determination of the enhancement of the enantiomeric purity during recrystallization of amino acids 139
Korhonen, I. O. O.
—— and Mäntykoski, K. M.
Gas–liquid chromatographic analyses. L. Retention, dispersion and selectivity indices of polychlorinated dibenzo-*p*-dioxins and dibenzofurans 327
Kraak, J. C., see Tock, P. P. H. 95
Kusz, P., see Szymanowski, J. 407
Lambert, W. J., see Dalga, R. J. 427
Larsson, L., see Drevin, I. 337
Lee, M. L., see Payne, K. M. 161
Lee, Y. W.
——, Cook, C. E., Fang, Q. C. and Ito, Y.
Application of analytical high-speed countercurrent chromatography to the isolation of bioactive natural products 434
Le Maréchal, P.
——, Decottignies, P., Jacquot, J.-P. and Miginiac-Maslow, M.
Separation by high-performance liquid chromatography of the ferredoxin–thioredoxin system proteins 305
Lottspeich, F., see Kehl, M. 131
Mabud, Md. A., see Hull, C. J. 387
McKinney, R. W., see Hull, C. J. 387
McNair, H. M., see Sandra, P. 63
Macrae, R., see Williams, R. A. 315
Maji, T., see Kawai, S. 467
Mäntykoski, K. M. 327
Maréchal, P. Le, see Le Maréchal, P. 305
Markides, K. E., see Payne, K. M. 161
Mertens, R., see Engelhardt, H. 169
Miginiac-Maslow, M., see Le Maréchal, P. 305
Mostert, I. A.
—— and Ramsteiner, K. A.
Coupled high-performance liquid chromatography–gas chromatography for the determination of pesticide residues in biological matrices 359

Müller, F.
——, Müller, H. and Straub, H.
Dosier- und Trennsäulenschaltsysteme für die hochauflösende Gasanalytik 25
Müller, H., see Müller, F. 25
Murayama, M., see Takeuchi, T. 147
Muschalek, V., see Koppenhoefer, B. 139
Nagano, H., see Kawai, S. 467
Nakazawa, H., see Ikai, Y. 397
Nishi, H.
——, Tsumagari, N., Kakimoto, T. and Terabe, S.
Separation of β-lactam antibiotics by micellar electrokinetic chromatography 259
Oka, H., see Ikai, Y. 397
Okamoto, Y.
——, Kaida, Y., Aburatani, R. and Hatada, K.
Optical resolution of amino acid derivatives by high-performance liquid chromatography on tris(phenylcarbamate)s of cellulose and amylose 367
Olsson, M.
——, Sander, L. C. and Wise, S. A.
Comparison of liquid chromatographic selectivity for polycyclic aromatic hydrocarbons on cyclodextrin and C_{18} bonded phases 277
Palamarev, H. E., see Palamareva, M. D. 235
Palamareva, M. D.
—— and Palamarev, H. E.
Microcomputer-aided characterization of mobile phases for normal-phase liquid–solid chromatography based on Snyder's theory and data 235
Pauthe-Dayde, D., see Henry, M. 413
Payne, K. M.
——, Davies, I. L., Bartle, K. D., Markides, K. E. and Lee, M. L.
Multidimensional packed capillary column supercritical-fluid chromatography using a flow-switching interface 161
Persson, B.-A., see Balmér, K. 107
Petersen, M., see Engelhardt, H. 169
Poppe, H., see Tock, P. P. H. 95
Prescott, S. R., see Hull, C. J. 387
Ramsey, R., see Katti, A. M. 119
Ramsteiner, K. A., see Mostert, I. A. 359
Rieder, R., see Kaiser, R. E. 49
Rijks, J., see Hetem, M. 3
——, see Van Es, A. 39
Roberts, D.
—— and Bertsch, W.
Practical aspects of recycle gas chromatography with capillary columns 59
Rochd, M., see Henry, M. 413
Rutten, G., see Hetem, M. 3
Sander, L. C., see Olsson, M. 277
Sandra, P.
——, David, F., Turner, K. A., McNair, H. M. and Brownstein, A. D.
Observations with high-molecular-weight polyethylene glycol stationary phases in capillary gas chromatography. I. Adsorption *versus* partitioning chromatography 63
Schill, G., see Balmér, K. 107
Schön, I., see Vadász, Z. 377
Seprödi, J., see Vadász, Z. 377
Shepherd, M. J., see Williams, R. A. 315
Ślebioda, M., see Wodecki, Z. J. 454
Snook, M. E.
——, Widstrom, N. W. and Gueldner, R. C.
Reversed-phase high-performance liquid chromatographic procedure for the determination of maysin in corn silks 439
Steenbeke, G., see Verzele, M. 87
Stein, H., see Hummel, M. 420
Straub, H., see Müller, F. 25
Suzuki, M., see Ikai, Y. 397
Szewczyk, H., see Szymanowski, J. 407
Szymanowski, J.
——, Kusz, P. and Szewczyk, H.
Increments of the arithmetic retention index for polyoxyethylene glycol monoalkyl ethers and their degradation products 407
Takeuchi, T.
——, Murayama, M. and Ishii, D.
Indirect photometric detection of cyclodextrins via inclusion complexation in micro high-performance liquid chromatography 147
Taylor, D. C., see Hull, C. J. 387
Teplán, I., see Vadász, Z. 377
Terabe, S., see Nishi, H. 259
Thiebaut, D.
——, Chervet, J.-P., Vannoort, R. W., De Jong, G. J., Brinkman, U. A. Th. and Frei, R. W.
Supercritical-fluid extraction of aqueous samples and on-line coupling to supercritical-fluid chromatography 151
Tock, P. P. H.
——, Boshoven, C., Poppe, H., Kraak, J. C. and Unger, K. K.
Performance of porous silica layers in open-tubular columns for liquid chromatography 95
Tsumagari, N., see Nishi, H. 259
Tun, L.
——, Wenzao, L. and Guoshi, T.
Separation and determination of 8β-hydroxyasterolid and perlolyrine in *Codonopsis pilosula* by reversed-phase high-performance liquid chromatography 458
Turner, K. A., see Sandra, P. 63
Unger, K. K., see Tock, P. P. H. 95

Vadász, Z.
——, Seprödi, J., Érchegyi, J., Teplán, I. and Schön, I.
High-performance liquid chromatographic monitoring of transpeptidation reactions in analogues of gonadotropin releasing hormone containing aspartic acid derivatives in position six 377
Van de Ven, L., see Hetem, M. 3
Van Es, A.
——, Rijks, J. and Cramers, C.
Turbulent flow in capillary gas chromatography 39
Vannoort, R. W., see Thiebaut, D. 151
Ven, L. van de, see Hetem, M. 3
Vermeer, B., see Hetem, M. 3
Verzele, M.
——, Steenbeke, G. and Vindevogel, J.
Micro-liquid chromatography with diode array detection 87
Vindevogel, J., see Verzele, M. 87
Vohra, R. M.
—— and Dube, S.
Identification and quantitation of rifamycins by reversed-phase high-performance liquid chromatography 463
Wenzao, L., see Tun, L. 458
Wetlaufer, D. B., see Gehas, J. 249
Widstrom, N. W., see Snook, M. E. 439
Williams, R. A.
——, Macrae, R. and Shepherd, M. J.
Non-aqueous size-exclusion chromatography coupled on-line to reversed-phase high-performance liquid chromatography. Interface development and applications to the analysis of low-molecular-weight contaminants and additives in foods 315
Wise, S. A., see Olsson, M. 277
Wodecki, Z. J.
——, Ślebioda, M. and Kołodziejczyk, A. M.
High-performance liquid chromatographic improvement of the Young racemization test 454
Yamada, M., see Ikai, Y. 397